

Hochschild two-cocycles and the good triple ($As, Hoch, Mag^\infty$)¹

Philippe LEROUX

Abstract: Hochschild two-cocycles play an important role in the deformation à la Gerstenhaber of associative algebras. The aim of this paper is to introduce the category of Hoch-algebras whose objects are associative algebras equipped with an extra magmatic operation $\succ$ verifying the Hochschild two-cocycle relation:

$$\mathcal{R}_2 : (x \succ y) * z + (x * y) \succ z = x \succ (y * z) + x * (y \succ z).$$

The free Hoch-algebra over a K -vector space is given in terms of planar rooted trees and the triples of operads $(As, Hoch, Mag^\infty)$ endowed with the infinitesimal relations are shown to be good. We then obtain an equivalence of categories between connected infinitesimal $Hoch$ -bialgebras and Mag^∞ -algebras.

Notation: In the sequel K is a field. We adopt Sweedler notation for the binary cooperation Δ on a K -vector space V and set $\Delta(x) = x_{(1)} \otimes x_{(2)}$. For a K -vector space V , we set $\bar{T}(V) := \bigoplus_{n \geq 0} V^{\otimes n}$.

1 Introduction

The well-known Poincaré-Birkhoff-Witt and the Cartier-Milnor-Moore theorems together can be rephrased as follows:

Theorem 1.1 (*CMM-PBW*) *For any cocommutative (associative) bialgebra $\mathcal{H}$, $Com^c - As$ -bialgebra for short, the following is equivalent.*

1. $\mathcal{H}$ is connected;

¹ 2000 Mathematics Subject Classification: 05E99, 16W30, 16W99, 18D50.

Key words and phrases: Hoch-algebras, infinitesimal Hoch-algebras, magmatic algebras, good triples of operads, cocycles d'Hochschild.

Email: ph_ler_math@yahoo.com

2. $\mathcal{H}$ is isomorphic to $U(\text{Prim } \mathcal{H})$ as a bialgebra;
3. $\mathcal{H}$ is isomorphic to $\text{Com}^c(\text{Prim } \mathcal{H})$ as a coalgebra,

where U is the usual enveloping functor and $\text{Prim } \mathcal{H}$ the usual Lie algebra of the primitive elements of $\mathcal{H}$.

In the theory developed by J.-L. Loday [3], this result is rephrased by saying that the triple of operads $(\text{Com}, \text{As}, \text{Lie})$, endowed with the usual Hopf relation, is good, where Com , As , and Lie stand respectively for the operads of commutative, associative and Lie algebras. Other good triples of operads equipped with other relations than the usual Hopf one, have been found since. A summary can be found in [3], see also [2, 1] for other examples.

It has been shown in [1] that the triple of operads $(\text{As}, \text{Dipt}, \text{Mag}^\infty)$ endowed with the semi-infinitesimal relations is good. The operad Dipt is related to dipterous algebras which are associative algebras equipped with an extra left module on themselves, see also [4], and Mag^∞ is related to Mag^∞ -algebras, i.e., K -vector spaces having one n -ary (magmatic) generating operation for each integer $n > 1$. We then obtained that the category of connected infinitesimal dipterous bialgebras, $\text{As}^c - \text{Dipt}$ -bialgebras for short, was equivalent to the category of Mag^∞ -algebras. In this paper, we propose another equivalence of category involving Mag^∞ : the category of connected infinitesimal *Hoch*-bialgebras is equivalent to the category of Mag^∞ -algebras.

In Section 2, we introduce *Hoch*-algebras and give an explicit construction of the free *Hoch*-algebra over a K -vector space. In Section 3, we introduce the notion of (connected) infinitesimal *Hoch*-bialgebras. In Section 4 we prove the announced equivalence of categories. In Section 5, we deal with unital *Hoch*-algebras and close by Section 6 with two other good triples involving the operad *Hoch*.

2 The free *Hoch*-algebra

A *Hoch*-algebra G is a K -vector space equipped with an associative operation $*$ and a magmatic operation $\succ$ verifying:

$$\mathcal{R}_2 : (x \succ y) * z + (x * y) \succ z = x \succ (y * z) + x * (y \succ z),$$

for all $x, y, z \in G$. Let V be a K -vector space. The free *Hoch*-algebra over V is defined as follows. It is equipped with a linear map $i : V \rightarrow \text{Hoch}(V)$ and for any *Hoch*-algebra G and any linear map $f : V \rightarrow G$, there exists a unique *Hoch*-algebra morphism $\phi : \text{Hoch}(V) \rightarrow G$ such that $\phi \circ i = f$. We now give an explicit construction of the free *Hoch*-algebra over a K -vector space.

Denote by T_n the set of rooted planar trees (degrees at least 2) with n leaves. The cardinalities of T_n are registered under the name *A001003 little Schroeder numbers* of the Online Encyclopedia of Integer Sequences. For $n = 1, 2, 3$, we get:

$$T_1 = \{ | \}, T_2 = \{ \vee \}, T_3 = \{ \vee\vee, \vee\vee, \vee\vee \}.$$

Define grafting operations by:

$$[\cdot, \dots, \cdot] : T_{n_1} \times \dots \times T_{n_p} \rightarrow T_{n_1 + \dots + n_p}, \quad (t_1, \dots, t_p) \mapsto [t_1, \dots, t_p] := t_1 \vee \dots \vee t_p,$$

where the tree $t_1 \vee \dots \vee t_p$ is the tree whose roots of the t_i have been glued together and a new root has been added. Observe that any rooted planar tree t can be decomposed in a unique way via the grafting operation as $t_1 \vee \dots \vee t_p$. Set $T_\infty := \bigoplus_{n \geq 0} K T_n$. Define over $\bar{T}(T_\infty)$, the following binary operations, first on trees, then by bilinearity:

$$\text{Concatenation} : (t_1 \dots t_p) * (s_1 \dots s_q) := t_1 \dots t_p s_1 \dots s_q,$$

$$(t_1 \dots t_p) \succ (s_1 \dots s_q) := \sum_{k=1}^q \sum_{i=0}^{p-1} t_1 \dots t_{p-(i+1)} [t_{p-i}, \dots, t_p, s_1, \dots, s_k] s_{k+1} \dots s_q.$$

For instance we get:

$$\begin{aligned} ||| \succ | &:= || \vee + | \vee\vee + \vee\vee \\ | \succ | \vee &:= \vee \vee + \vee\vee \end{aligned}$$

Theorem 2.1 *The K -vector space $\bar{T}(T_\infty)$ endowed with the operations $*$ and $\succ$ is the free Hoch-algebra over K .*

Proof: Let $x := x_1 \dots x_m$, $y := y_1 \dots y_n$ and $z := z_1 \dots z_p$. We get:

$$\begin{aligned} (x \succ y) * z + (x * y) \succ z &= \sum_{k=1}^n \sum_{i=0}^{m-1} x_1 \dots x_{m-(i+1)} [x_{m-i}, \dots, x_m, y_1, \dots, y_k] y_{k+1} \dots y_n z_1 \dots z_p \\ &+ \sum_{k=1}^p \sum_{i=0}^{n-1} x_1 \dots x_m y_1 \dots y_{n-(i+1)} [y_{n-i}, \dots, y_n, z_1, \dots, z_k] z_{k+1} \dots z_p \\ &+ \sum_{k=1}^p \sum_{i=0}^{m-1} x_1 \dots x_{m-(i+1)} [x_{m-i}, \dots, x_m, y_1, \dots, y_n, z_1, \dots, z_k] z_{k+1} \dots z_p \\ \\ x \succ (y * z) + x * (y \succ z) &= \sum_{k=1}^n \sum_{i=0}^{m-1} x_1 \dots x_{m-(i+1)} [x_{m-i}, \dots, x_m, y_1, \dots, y_k] y_{k+1} \dots y_n z_1 \dots z_p \\ &+ \sum_{k=1}^p \sum_{i=0}^{m-1} x_1 \dots x_{m-(i+1)} [x_{m-i}, \dots, x_m, y_1, \dots, y_n, z_1, \dots, z_k] z_{k+1} \dots z_p \\ &+ \sum_{k=1}^p \sum_{i=0}^{n-1} x_1 \dots x_m y_1 \dots y_{n-(i+1)} [y_{n-i}, \dots, y_n, z_1, \dots, z_k] z_{k+1} \dots z_p, \end{aligned}$$

showing that

$$(x \succ y) * z + (x * y) \succ z = x \succ (y * z) + x * (y \succ z),$$

holds for all forests of planar rooted trees x, y, z . Observe that any rooted planar tree $t := [t_1, \dots, t_n]$ can be rewritten as:

$$t = (t_1 * (t_2 \dots t_{n-1})) \succ t_n - t_1 * ((t_2 \dots t_{n-1}) \succ t_n)$$

Let G be a *Hoch*-algebra and $g \in G$ and $f : K \rightarrow G$ be a linear map. Consider the embedding $i : K \hookrightarrow \bar{T}(T_\infty)$ defined by $i(1_K) := |$ and define by induction the map $\phi : \bar{T}(T_\infty) \rightarrow G$ as follows:

$$\phi(|) = g,$$

$$\phi(t_1 \dots t_n) = \phi(t_1) *_G \phi(t_2) *_G \dots *_G \phi(t_n),$$

$$\phi(t) = (\phi(t_1) *_G (\phi(t_2) \dots \phi(t_{n-1}))) \succ_G \phi(t_n) - \phi(t_1) *_G ((\phi(t_2) \dots \phi(t_{n-1})) \succ_G \phi(t_n))$$

for any $t := [t_1, \dots, t_n]$ and extend ϕ by linearity. By construction, ϕ is a morphism of associative algebras. Using the fact that:

$$(x * y) \succ z - x * (y \succ z) = x \succ (y * z) - (x \succ y) * z,$$

and changes of indices in the involving sums, one shows that ϕ is also a morphism for the magmatic operations. It is then the only *Hoch*-morphism such that $\phi \circ i = f$. $\square$

As the operad *Hoch* is regular, the following holds. Let V be a K -vector space. The free *Hoch*-algebra over V is the K -vector space:

$$Hoch(V) := \bigoplus_{n>0} Hoch_n \otimes V^{\otimes n},$$

with $Hoch(K) := \bigoplus_{n>0} Hoch_n \simeq \bar{T}(T_\infty)$ (hence $Hoch_n$ is explicitly described in terms of forests of rooted planar trees) equipped with the operations $*$ and $\succ$ defined as follows:

$$((t_1 \dots t_n) \otimes \omega) * ((s_1 \dots s_p) \otimes \omega') = (t_1 \dots t_n s_1 \dots s_p) \otimes \omega \omega',$$

$$((t_1 \dots t_n) \otimes \omega) \succ ((s_1 \dots s_p) \otimes \omega') = ((t_1 \dots t_n) \succ (s_1 \dots s_p)) \otimes \omega \omega',$$

for any $\omega \in V^{\otimes n}, \omega' \in V^{\otimes p}$. The embedding map $i : V \hookrightarrow Hoch(V)$ is defined by: $v \mapsto | \otimes v$.

Since the generating function associated with the Schur functor $\bar{T}$ is $f_{\bar{T}}(x) := \frac{x}{1-x}$ and with the Schur functor T_∞ is $f_{T_\infty}(x) := \frac{1+x-\sqrt{1-6x+x^2}}{4} = x + x^2 + 3x^3 + 11x^4 + 45x^5 + \dots$, the generating function of the operad *Hoch* is $f_{\bar{T}} \circ f_{T_\infty}$, that is:

$$f_{Hoch}(x) := \frac{1+x-\sqrt{1-6x+x^2}}{3-x+\sqrt{1-6x+x^2}} = x + 2x^2 + 6x^3 + 22x^4 + \dots$$

The sequence $(1, 2, 6, 22, 90, \dots)$ is registered as A006318 under the name *Large Schroeder numbers* on the Online Encyclopedia of Integer Sequences.

3 Infinitesimal Hoch-bialgebras

By definition, an infinitesimal Hoch-bialgebra (or an As^c – Hoch-bialgebra for short) $(\mathcal{H}, *, \succ, \Delta)$ is a Hoch-algebra equipped with a coassociative coproduct Δ verifying the following so-called nonunital infinitesimal relations:

$$\Delta(x \succ y) := x_{(1)} \otimes (x_{(2)} \succ y) + (x \succ y_{(1)}) \otimes y_{(2)} + x \otimes y.$$

$$\Delta(x * y) := x_{(1)} \otimes (x_{(2)} * y) + (x * y_{(1)}) \otimes y_{(2)} + x \otimes y.$$

It is said to be connected when $\mathcal{H} = \bigcup_{r \geq 0} F_r \mathcal{H}$ with the filtration $(F_r \mathcal{H})_{r \geq 0}$ defined as follows:

$$(\text{The primitive elements}) \quad F_1 \mathcal{H} := \text{Prim } \mathcal{H} = \ker \Delta,$$

Set $\Delta^{(1)} := \Delta$ and $\Delta^{(n)} := (\Delta \otimes id_{n-1}) \Delta^{(n-1)}$ with $id_{n-1} = \underbrace{id \otimes \dots \otimes id}_{\text{times } n-1}$. Then,

$$F_r \mathcal{H} := \ker \Delta^{(r)}.$$

Theorem 3.1 *Let V be a K -vector space. Define on $\text{Hoch}(V)$, the free Hoch-algebra over V , the cooperation $\Delta : \text{Hoch}(V) \rightarrow \text{Hoch}(V) \otimes \text{Hoch}(V)$ recursively as follows:*

$$\Delta(i(v)) := 0, \text{ for all } v \in V,$$

$$\Delta(x \succ y) := x_{(1)} \otimes (x_{(2)} \succ y) + (x \succ y_{(1)}) \otimes y_{(2)} + x \otimes y.$$

$$\Delta(x \star y) := x_{(1)} \otimes (x_{(2)} \star y) + (x \star y_{(1)}) \otimes y_{(2)} + x \otimes y,$$

for all $x, y \in \text{Hoch}(V)$. Then $(\text{Hoch}(V), \Delta)$ is a connected infinitesimal Hoch-bialgebra.

Proof: This result can be proved by hand or can be seen as a corollary of the Theorem 4.2 in the next section. $\square$

4 A good triple of operads

It can be usefull to have the following result when searching for good triples.

Lemma 4.1 *Let $\mathcal{C}, \mathcal{A}, \mathcal{Z}, \mathcal{Q}$ and Prim be operads. Suppose the triples of operads $(\mathcal{C}, \mathcal{A}, \text{Prim})$ and $(\mathcal{C}, \mathcal{Z}, \text{Vect})$ equipped with the same compatibility relations to be good. Suppose $\mathcal{A} = \mathcal{Z} \circ \mathcal{Q}$ then $\text{Prim} = \mathcal{Q}$.*

Proof: Since $(\mathcal{C}, \mathcal{Z}, \text{Vect})$ is good, the notion of $\mathcal{C}^c$ – $\mathcal{Z}$ -bialgebra has a meaning and the following is equivalent:

1. The $\mathcal{C}^c - \mathcal{Z}$ -bialgebra $\mathcal{H}$ is connected.
2. As $\mathcal{Z}$ -algebras, $\mathcal{H}$ is isomorphic to the free $\mathcal{Z}$ -algebra over its primitive elements.
3. As $\mathcal{C}^c$ -coalgebras, $\mathcal{H}$ is isomorphic to the cofree $\mathcal{C}^c$ -coalgebra over its primitive elements.

As $(\mathcal{C}, \mathcal{A}, Prim)$ is good, the isomorphism of Schur functors $\mathcal{A} \simeq \mathcal{C}^c \circ Prim$ holds. Therefore, if V is a K -vector space, then $\mathcal{A}(V) = \mathcal{C}^c(Prim(V))$ and by hypothesis $\mathcal{A}(V) = \mathcal{Z}(\mathcal{Q}(V))$. Hence $\mathcal{A}(V)$ is a connected $\mathcal{C}^c - \mathcal{Z}$ -bialgebra. Consequently, $Prim = \mathcal{Q}$. $\square$

Theorem 4.2 *The triple of operads $(As, Hoch, Mag^\infty)$ endowed with the infinitesimal relation is good.*

Proof: Fix an integer $n > 0$. By $[n] - Mag$ we mean the regular binary operad generated by n magmatic (binary) operations. In [2] Thm. 4.4 (and 4.5), it has been shown that for each integer $n > 0$ the triples of operads $(As, [n] - Mag, Prim [n] - Mag)$ endowed with the infinitesimal relations were good. For $n = 2$, the operadic ideal J generated by the primitive operations:

$$\begin{aligned} &*(\succ \otimes id) + \succ (* \otimes id) - \succ (id \otimes *) - *(id \otimes \succ), \\ &*(\ast \otimes id) - *(id \otimes \ast), \end{aligned}$$

yields another good triple of operads $(As, [2] - Mag/J, Prim ([2] - Mag/J))$ (cf. Prop 3.1.1 [3] on quotient triples), which turns out to be the triple $(As, Hoch, Prim Hoch)$. As $(As, As, Vect)$ endowed with the infinitesimal relation is good (cf [4]) and $Hoch = As \circ Mag^\infty$ using Section 2, we get $Prim Hoch = Mag^\infty$ by using Lemma 4.1. $\square$

We then obtain another equivalence of categories involving the operad Mag^∞ .

Corollary 4.3 *The category of connected infinitesimal $As^c - Hoch$ -bialgebras and the category of Mag^∞ -algebras are equivalent.*

$$\{conn. As^c - Hoch - bialg.\} \xrightleftharpoons[Primitive]{U} \{Mag^\infty - alg.\},$$

where U and $Primitive$ are respectively the universal enveloping functor and the primitive functor.

Proof: Apply Thm. 2.6.3. [3]. $\square$

Remark: The functor $Primitive$ is obviously given as follows. If $(\mathcal{H}, *, \succ)$ is a connected infinitesimal $Hoch$ -bialgebra, then for all integer $n > 1$ and for all primitive elements $x_1, \dots, x_n \in \mathcal{H}$, the element:

$$[x_1, \dots, x_n]_n := (x_1 * \dots * x_{n-1}) \succ x_n - x_1 * ((x_2 * \dots * x_{n-1}) \succ x_n),$$

will be primitive. The functor U acts as follows. Let $(M, ([\dots, \cdot]_n)_{n \geq 1})$ be a Mag^∞ -algebra with the $[\dots, \cdot]_n$ being its generating n -ary operations. Then $U(M)$ is given by $Hoch(M)/\sim$, where the equivalence relation $\sim$ consists in identifying,

$$(x_1 * \dots * x_{n-1}) \succ x_n - x_1 * ((x_2 * \dots * x_{n-1}) \succ x_n),$$

with $[x_1, \dots, x_n]_n$, for all $x_1, \dots, x_n \in M$.

5 Extension to a unit

Unital $Hoch$ -algebras are $Hoch$ -algebras equipped with a unit 1 whose compatibility with operations are defined as follows:

$$x \succ 1 = x = 1 \succ x, \quad x * 1 = x = 1 * x.$$

For instance, $Hoch_+(V) := K \cdot 1_K \oplus Hoch(V)$, where $Hoch(V)$ is the free $Hoch$ -algebra over a K -vector space V is a unital $Hoch$ -algebra with unit 1_K . This gives birth to unital Mag^∞ -algebras which are Mag^∞ -algebras such that the generating operations are related with the unit as follows:

$$\begin{aligned} [1, \cdot, \dots, \cdot]_n &= 0, \\ [\cdot, \dots, 1, \dots, \cdot]_n &= [\cdot, \dots, \cdot]_{n-1}, \\ [\cdot, \dots, \cdot, 1]_n &= 0. \end{aligned}$$

Over $Hoch_+(V)$, one has a unital infinitesimal coproduct δ defined via the former coproduct Δ as follows:

$$\delta(x) = 1_K \otimes x + x \otimes 1_K + \Delta(x),$$

for any $x \in Hoch(V)$. The compatibility relations are the so-called unital infinitesimal relations defined as follows:

$$\begin{aligned} \Delta(x \succ y) &:= x_{(1)} \otimes (x_{(2)} \succ y) + (x \succ y_{(1)}) \otimes y_{(2)} - x \otimes y. \\ \Delta(x * y) &:= x_{(1)} \otimes (x_{(2)} * y) + (x * y_{(1)}) \otimes y_{(2)} - x \otimes y. \end{aligned}$$

We then obtain the good triple of operads $(As, Hoch, Mag^\infty)$ equipped with the unital infinitesimal relations.

6 Other triples of operads

The triple of operads $(As, Hoch, Mag^\infty)$ endowed with the infinitesimal relations are not the only one involving the operad $Hoch$. By changing the compatibility relations, two other good triples of operads $(Com, Hoch, Prim_{Com} Hoch)$ and $(As, Hoch, Prim_{As} Hoch)$ endowed respectively with the Hopf relations and the semi-Hopf relations can be proposed. But contrary to the case of the triple $(As, Hoch, Mag^\infty)$ the explicit descriptions of operads of the primitive elements of these two other triples are open problems.

References

- [1] Ph. LEROUX. Infinitesimal or cocommutative dipterous bialgebras and good triples of operads. *arXiv:0803.1421*.
- [2] Ph. LEROUX. L-algebras, triplicial-algebras, within an equivalence of categories motivated by graphs. *arXiv:0709.3453v2*.
- [3] J.-L. LODAY. Generalized bialgebras and triples of operads. *arXiv:math.QA/0611885*.
- [4] J.-L. LODAY and M. RONCO. On the structure of cofree Hopf algebras. *J. reine angew. Math.*, 592:123–155, 2006.